

Quantitative Evaluation Methods for Secondary Ecosystem Management Activities -For *Satoyama* Banking-

Akira TANAKA, Takafumi KAWAMURA
Tokyo City University

1 Background and Purpose

A primary ecosystem has already disappeared in many countries, and much of the nature that remains is a secondary ecosystem managed with traditional human activities. In Japan, these secondary ecosystems are called “*Satoyama*”. Traditionally wisely used *Satoyama* maintains biodiversity that is different from, but not inferior to, primary ecosystem.

However, secondary ecosystems, such as *Satoyama* in Japan, continue to decrease due to development projects. And even if they remain, they face biodiversity degradation due to a lack of traditional management. The former problem is called “overuse” in terms of “space,” and the latter is called “underuse” in terms of “quality.”

To address the overuse problem, assessment of “no net loss” has been introduced by comparing the negative impact on nature and the positive impact of compensatory mitigation such as biodiversity offsets. Such assessment is based on quantitative habitat assessments represented by HEP, Habitat Evaluation Procedure, which is part of the environmental impact assessment.

In this study, we devised a quantitative evaluation method based on HEP. Then, it was demonstrated with a case of secondary ecosystem management activities by people including a non-profit organization. The results were compared with an overuse case of a land development project, for which a quantitative evaluation was performed in the EIA. Based on this comparative analysis, we propose requirements for an effective quantitative assessment method to evaluate the effects on secondary ecosystem management activities in future.

2 Methodology

2.1 Quantitative assessment of the impacts of management activities in secondary ecosystems

We evaluated the ecosystem management activities in Aichi Prefecture, using *Luehdorfia japonica* as an indicator

species of the *Satoyama* ecosystem. For the evaluation, SIs (Suitability Indices) and HSI (Habitat Suitability Index) models were constructed and calculated. The management outcomes were assessed by comparing the THU (Total Habitat Unit) before and after the management activities.

2.2 Quantitative assessment of the impacts of development projects in secondary ecosystems

We evaluated the impact of a land development project in a *Satoyama* in Yokohama City using two species of fireflies (*Luciola cruciata* and *Luciola lateralis*) and two species of frogs (*Rana ornativentris* and *Rana japonica*) as indicators of the *Satoyama* ecosystem including wetlands. SI and HSI models were constructed from previous studies of these species and calculated. The development impacts were assessed by comparing THUs before development and THUs for three development scenarios, including no-action.

2.3 Issues in quantitative assessment of the impacts of management activities in secondary ecosystems

A comparison between the quantitative assessment of the impacts of management activities and development projects in secondary ecosystems revealed issues of consideration in the quantitative assessment of the impacts of management activities in secondary ecosystems such as *Satoyama* ecosystems.

The points of comparison were the type of human activity, the type of *Satoyama* ecosystem, the purpose of the assessment, the species used in the assessment, the method of determining the assessment area, and the assessment results. And it is necessary tool to assess the quality of secondary ecosystems by assessing effects of people’s management activities in biodiversity offset in secondary ecosystems.

3. Results

3.1 Quantitative assessment of the impacts of management activities in secondary ecosystems

Habitat variables were selected as the availability of plants whose flowering season is from March to May, the availability of space for flight, and the availability of *Asarum* spp.

For the evaluation area, habitat evaluation was conducted using a 25m square evaluation grid.

Figure 1 shows the relationship between the habitat variables, SI models, and HSI model in the adult and larval stages of the *L. japonica*.

The respective SI models before and after the management activities in the experimental site are shown in Tables 1 to 4, and the HSI model is shown in Equation 1. Figures 2 to 7 show the respective evaluation results, and Figure 8 shows the graph of THU before and after the management activities.

For the SI2 model on the availability of flying space, two separate models (SI2 α and SI2 β) were developed to account for the consideration of spatial connectivity.

Some grids assessed in this study included areas where no management activities were conducted, and in such areas, the HSI values were lower than in areas where activities were conducted. Furthermore, the SI2 β model, which took into consideration spatial connectivity, yielded a higher SI value than the SI2 α model, even in the areas where management activities are not conducted in the surrounding grids. However, since the HSI model was constructed to change significantly depending on the availability of *Asarum* spp., the overall HSI value became low even when the SI2 β value was high.

$$HSI = \frac{SI1 + SI2}{2} \times SI3$$

Equation 1 HSI model for evaluating the outcomes of Satoyama management activities




Figure 1 Relationship between the Habitat Variables, SI models, and HSI model of the *L. japonica*

Table 1 SI model for SI1 (the availability of plants whose flowering season is from March to May)

Condition	SI value
Have not seen any plants with purple or yellow flowers within a year.	0.0
Have identified plants with yellow flowers within a year.	0.5
Have identified plants with purple flowers within a year.	1.0

Table 2 SI model for SI2 α (the availability of space for flight, without consideration of connectivity)

Condition	SI value
Type A (very high canopy density and forest density, thickets on the forest floor)	0.0
Type B (density with moderate sunlight in the forest)	0.5
Type C (very low canopy density and forest density, forest floor is not thicketed)	1.0

Table 3 SI model for SI2 β (the availability of space for flight, with consideration of connectivity)

Condition	SI value
Evaluation grid is Type A and all surrounding 8 grids are also Type A.	0.0
Evaluation grid is Type A and any of the surrounding 8 grids contain Type B or C.	0.2
Evaluation grid is Type B and all surrounding 8 grids are also Type B.	1.0
Evaluation grid is Type B and any of the surrounding 8 grids contain Type A or C.	0.5
Evaluation grid is Type C and all surrounding 8 grids are also Type C.	0.0
Evaluation grid is Type C and any of the surrounding 8 grids contain Type A and B.	0.8

Table 4 SI model for SI3 (the availability of *Asarum* spp)

Condition	SI value
No growth of <i>Asarum rigescens</i> var. <i>brachypodium</i> or <i>Asarum takaoi</i> .	0.5
Growth of <i>Asarum rigescens</i> var. <i>brachypodium</i> or <i>Asarum takaoi</i> can be seen.	1.0




Figure 2 SI1 values before and after management activities



Figure 3 SI2α values before and after management activities




Figure 4 SI2β values before and after management activities




Figure 5 SI3 values before and after management activities




Figure 6 HSI values before and after management activities when SI2α model is used




Figure 7 HSI values before and after management activities with SI2β model is used




Figure 8 THU before and after management activities

3.2 Quantitative assessment of the impacts of development projects in secondary ecosystems

The habitat assessment results of the four species are shown in Figures 9 to 12. Except for *R. ornativentris*, THU decreased in all scenarios, including no-action.




Figure 9 Change in THU of *Luciola cruciata*




Figure 10 Change in THU of *Luciola lateralis*




Figure 11 Change in THU of *Rana ornativentris*




Figure 12 Change in THU of *Rana japonica*

4. Discussion

The results of the two case studies are summarized in Table 5 according to the perspectives on quantitative biodiversity assessment.

We identified five issues to consider when developing a quantitative evaluation method for the impacts of secondary ecosystem management activities.

First, the method should be designed to be utilized by those who do not have expertise in biology. Since HEP is a quantitative biological assessment, which is time-consuming and costly, it needs to be conducted by experts in biology and ecology. It requires detailed and diverse data for decision-making, including a selection of target species, construction of SI and HSI models, and estimation of SI values. To effectively conserve *Satoyama* ecosystems, it is necessary to develop a simple assessment method that can be utilized by non-specialists in biology and ecology.

Second, the method should convey the causal relationship between specific management activities and the conservation impacts on ecosystems and habitats. HEP is a quantitative biological assessment and cannot directly evaluate the impacts of human management activities. However, in the case of species in *Satoyama* ecosystems whose habitats have been maintained through human involvement, the management activities themselves are directly related to the quality of the habitat.

Third, the method should evaluate human management activities themselves since HEP evaluates the biological status of the target area as a habitat and not the management activities directly.

Fourth, the method should evaluate the impacts of continuous management activities. HEP evaluates the biological status of a target area at a certain point in time. In constant human management activities, such as in *Satoyama* ecosystems, extensive research efforts become necessary to effectively evaluate the changes over time.

Lastly, the method should be simple and yet the ecological health aspect of HEP should not be compromised.

The results of HEP are quantitative evaluations from an ecological perspective. Therefore, it may not be easy to translate the value of ecosystems and incorporate it into the cost of activities in *Satoyama* ecosystem management planning and trade-offs involving biodiversity offsetting and banking in the future. It would be possible to better integrate the benefits of conservation activities holistically through developing a method to evaluate the management activities themselves in a simple and quantitative manner without compromising the perspective of evaluating the ecological health of the HEP.

Table 5 Perspectives on quantitative biodiversity assessment and the results of the case studies

Perspectives	3.1. Yokohama Project	3.2. Aichi Project
Type of human activities	Residential development business	Satoyama conservation activities
Type of <i>Satoyama</i> ecosystem (Area)	Deciduous broad-leaved forest, paddy field (33.6ha)	Deciduous broad-leaved forest (4.0ha)
Evaluation target	Impacts of development project and mitigation measures on the habitat of evaluation species	Impacts of conservation activities on the habitat of evaluation species
Evaluation species used for HEP	Genji firefly (<i>Luciola cruciate</i>), Heike firefly (<i>Luciola lateralis</i>), Japanese brown frog (<i>Rana japonica</i>), Montane brown frog (<i>Rana ornativentris</i>)	Japanese luehdorfia (<i>Luehdorfia japonica</i>)
Evaluation area division method	Polygon	Mesh
Evaluation results	Negative impacts of the development project could be evaluated biologically and quantitatively for each evaluation species	Positive impacts of outcomes of the conservation activities could be evaluated biologically and quantitatively for each evaluation species

Reference

- Tanaka A (2010) “SATOYAMA Banking” to Solve Satoyama’s Overuse and Underuse Problems: Integration of Biodiversity Banking, Strategic Environmental Assessment and Satoyama Conservation (Japanese language). p47-51, Council of Environmental local government, Environmental Autonomy White Paper 2010 Edition. Seikatsusha, Tokyo, 180pp.
- Kuki N, Tanaka A, Makita K, Ogura S, Ito Y (2011) Development of “Easy HEP,” a biodiversity assessment method using the HSI Chart. Japan Society for Environmental Assessment 2011, p162-167.
- Uchiike T, Matsumiya A, Tanaka A (2017) A Study on Biodiversity Assessment Methods in Urban Areas. Architectural Institute of Japan 2017, p937-938.
- Watabe Y, Ide K, Saiki S, Aoshima M, Shiromoto T, Togo Y, Hori Y, Matsuoka A, Watanabe T, Oda S, Tanaka A (2017) A Study on the Challenges of Applying Overseas Valuation Methods to Biodiversity Offsetting in Satoyama. Japan Society for Environmental Assessment 2017, p141-144.
- Shirouzu T, Kawazoe A, Wakamatsu M (1976) Original Color Japanese Butterfly Catalogue. Hoikusha, 422pp.
- Fujisawa S (1983) *Luehdorfia japonica* and Asarum. *Luehdorfia japonica* research group. 402pp.
- Aichi Prefecture (2016) *Luehdorfia japonica* Conservation Plan. Nature Environment Division, Environment Department, Aichi Prefecture, 33pp.
- Aichi Prefecture (2016) Report on the Maritime Forest Survey No. 5, 2015. Aichi Maritime Forest Center, 205pp.
- Aichi Prefecture (2017a) Mitigation efforts at Expo 2005 Aichi Japan. <http://www.pref.aichi.jp/kankyo/sizen-ka/shizen/ecologicalnetwork/aichimitigation/kaisyonomori.pdf> (2017.09.19).
- Aichi Prefecture (2017b) Red List of Aichi Prefecture 2009. http://www.pref.aichi.jp/kankyo/sizen-ka/shizen/yasei/rdb/pdf/ANIMALS/ANIMALS08_2.pdf, (2017.09.19).
- Aoyama J (2005) Science Album *Luehdorfia japonica*. Akane Shobo Co., 54pp.
- Ozeki M, Suka T, Hamada T, Kishimoto R (2007) Selection of spawning environment for *Luehdorfia japonica*, Nagano Prefectural Institute for Environmental Conservation. p21-25.

- Kameyama T, Watanabe K (2011) Conservation and changes of flight behavior in *Luehdorfia japonica* (Papilionidae, Lepidoptera) at the Ege-san Municipal Park in Hiroshima City: Significance of a huge TV-tower, Hiroshima University Museum, p37-22.
- Kurita S, Oda K, Abe Y (2016) Home of *Luehdorfia japonica* and *Luehdorfia puziloi*, their mixed habitat and life history. Publication Committee of *Luehdorfia japonica*. Nagano Prefecture, Creative center, 160pp.
- The Japan Association for the International Exposition 2005 (1999) Environmental Impact Assessment for the 2005 Japan International Exposition.
- Shirouzu T (2006) Standard Catalogue of Butterflies from Japan. Gakushu-kenkyusha, 336pp.
- Tanaka A (2013) Introduction to HEP - "Habitat Evaluation Procedures" Manual - Theory and practices for Habitat Evaluation Procedure (HEP) in Japan. Asakura-Shoten, 280pp.
- Nakanishi M (1979) Food habits of *Luehdorfia japonica* on *Asarum* spp. I – VI.
- Natsuaki M, Takeuchi T (1999) Study on the movements of adult *Luehdorfia japonica* Leech (Lepidoptera, Papilionidae) by the mark-release-recapture method. Lepidoptera Science, Trans. lipid. Soc. Japan 50 (3); p216-222.
- Hamano E (1986) Nature and Science 56 *Luehdorfia japonica* - fly into spring -. Iwasaki-Shoten Co., 39pp.
- Hara S (1979) Natural history of *Luehdorfia japonica*. Tsukiji-shoten Co., 210pp.
- Hiroki S, Eda N, Hatta K (1999a) Distribution pattern of *Asarum rigescens* var. *brachypodion* and spawning density of *Luehdorfia japonica* in the Maritime Forest (Seto City, Aichi Prefecture, Japan). Association for Information Culture, No.9, p23-30.
- Hiroki S, Eda N, Hatta K (1999b) Population structure of *Asarum rigescens* var. *brachypodion* and spawning density of *Luehdorfia japonica* in the secondary forest of Maritime Forest (Seto City, Aichi Prefecture, Japan) in spring 1999. Association for Information Culture, No.10, p37-47.
- Mt. Monomi Nature Observation Meeting, Maritime Forest Network (1997) Handbook of Flowers, Insects and Trees in the Maritime Forest. Kobushi-shobo, 207pp.
- Watanabe Y (1996) *Luehdorfia japonica* Monograph of Luehdorfia Butterflies. Hokkaido University Library, 269pp.
- Katou (2016) Flora of Mikawa. <http://mikawanoyasou.org/>, (2016.1.15).
- Masuda M a (2016) Professor, Nagoya Institute of Technology, Interview by mail. 2016.11.2.
- Masuda M b (2016) Professor, Nagoya Institute of Technology, Interview by mail, 2016.11.16.
- Masuda M c (2016) Professor, Nagoya Institute of Technology, Interview by mail, Aichi Marine Forest. 2016.11.29.
- Masuda M (2017) Professor, Nagoya Institute of Technology, Interview by mail, 2017.2.6.
- Takahashi A (2017) Member of the Medical History Society of Japan, Interview. National Olympics Memorial Youth Center. 2017.2.11.
- Takahashi M (2016) Representative of the Kasugai *Luehdorfia japonica* Satoyama Creation Group, Interview, Aichi Marine Forest. 2016.11.29.
- Takahashi M b (2017) Representative of the Kasugai *Luehdorfia japonica* Satoyama Creation Group, Interview by mail. 2017.2.8.
- Takahashi Mayumi (2017) Member of the Lepidopterological Society, Interview. National Olympics Memorial Youth Center. 2017.2.11.
- Ishii T (2016) *Luehdorfia japonica* Research Group, Interview by mail, 2016.11.26.

End of Paper